

# HYPOGLYCAEMIA (Hypo)

**LOW**

Blood Glucose Level (BGL) < 4.0 mmol/L

**DO NOT LEAVE STUDENT ALONE  
DO NOT DELAY TREATMENT**

## Signs and symptoms

Note: Symptoms may not always be obvious.  
Pale, headache, shaky, sweaty, dizzy, drowsy, changes in behaviour.

**Student conscious**  
(Able to eat hypo food)

**Student unconscious/  
drowsy**  
(Risk of choking/unable to swallow)

**Step 1: Give fast acting carbohydrate**  
(As supplied or listed on management plan)

**First Aid DRSABCD**  
Stay with unconscious student

**Step 2: Recheck BGL after 15 mins**  
If BGL < 4.0 repeat Step 1  
If BGL ≥ 4.0 go to Step 3

**CALL AN  
AMBULANCE  
DIAL 000**

**Step 3: Give sustaining carbohydrate**  
(As supplied or listed on management plan)

**Contact parent/carer**  
When safe to do so

**PARENT/CARER NAME** \_\_\_\_\_

**CONTACT No** \_\_\_\_\_

**OTHER CONTACT NAME & No** \_\_\_\_\_

## SCHOOL SETTING

Multiple daily injections

# DIABETES ACTION PLAN 2018



**STUDENT'S NAME**

\_\_\_\_\_

**SCHOOL**

\_\_\_\_\_

Use this plan in conjunction with Diabetes Management Plan

- Continuous Glucose Monitoring (CGM) Appendix
- Flash Glucose Monitoring Appendix

## INSULIN

Insulin is given 4 or more times per day.

An injection will be needed before lunch.

- Able to inject insulin:
- independently
  - with supervision
  - with assistance

Injection will be given in: \_\_\_\_\_  
(Room/location)

## ROUTINE BGL CHECKING TIMES

- Anytime, anywhere in the school
- Before food
- Anytime hypo is suspected
- Before activity
- Before exams or tests (e.g. NAPLAN)

## PHYSICAL ACTIVITY

- 15 grams sustaining carbohydrate before every 30 mins of planned activity
- Vigorous activity should **not** be undertaken if BGL ≥ 15.0 and blood ketones are ≥ 1.0

# HYPERGLYCAEMIA (Hyper)

**HIGH**

Blood Glucose Level (BGL) ≥ 15.0 mmol/L

**HIGH BGLS ARE COMMON**

## Signs and Symptoms

Note: Symptoms may not always be obvious.  
Some could be increased thirst, extra toilet visits, poor concentration, irritability, tiredness

**Student well**  
Re-check BGL in 2 hours

**Student unwell**  
Check blood ketones (if able) if ≥ 1.0 mmol/L

**Encourage oral fluids, return to class**  
1-2 glasses water per hour; extra toilet visits may be required

**CONTACT PARENT/  
CARER  
TO COLLECT  
STUDENT ASAP**

**In 2 hours, if BGL still ≥ 15.0  
check ketones  
call parent/carer for advice**

**DATE** \_\_\_\_\_

**TREATING MEDICAL TEAM** \_\_\_\_\_

**CONTACT No** \_\_\_\_\_

